

On Ky Fan's Section Theorem*

Xian Wu and Feng Li

*Department of Mathematics, Yunnan Normal University, Kunming, Yunnan 650092,
People's Republic of China*

metadata, citation and similar papers at core.ac.uk

In this paper we first give a new section theorem of Ky Fan type and its equivalent theorem in an H -space, and next as their applications we obtain some minimax theorems and a coincidence theorem. © 1998 Academic Press

Key Words: H -space; H -convex set; upper (lower) semicontinuous function; section theorem; multivalued mapping.

1. INTRODUCTION AND PRELIMINARIES

In 1961, Ky Fan [3] proved the following section theorem:

THEOREM A. *Let X be a nonempty compact convex subset of a Hausdorff topological vector space E , and let A be a nonempty subset of $X \times X$ with $(x, x) \in A$ for all $x \in X$. Suppose the following conditions are fulfilled:*

- (i) *For each $x \in X$, the set $\{y \in X: (x, y) \notin A\}$ is convex or empty.*
- (ii) *For each $y \in X$, the set $\{x \in X: (x, y) \in A\}$ is closed.*

Then there exists a point $x_0 \in X$ such that $\{x_0\} \times X \subset A$.

In 1980, Ha [5] generalized Theorem A as follows:

THEOREM B. *Let E and F be Hausdorff topological vector spaces, let $X \subset E$, $Y \subset F$ be nonempty convex subsets, and let A be a subset of $X \times Y$ such that*

- (i) *For each $x \in X$, the set $\{y \in Y: (x, y) \notin A\}$ is convex or empty.*
- (ii) *For each $y \in Y$, the set $\{x \in X: (x, y) \in A\}$ is closed in X .*

* Sponsored by the National Science Foundation of China and the Foundation of the Yunnan Science and Technology Commission and the Foundation of the Yunnan Educational Commission.

Suppose that there exists a subset B of A and a compact convex subset K of X such that B is closed in $X \times Y$ and

(iii) For each $y \in Y$, the set $\{x \in K: (x, y) \in B\}$ is nonempty and convex. Then there exists a point $x_0 \in K$ such that $\{x_0\} \times Y \subset A$.

This paper has two purposes. First we give a section theorem of the Ky Fan type and its equivalent theorem in an H -space, and next, as their applications, we obtain some minimax theorems and some coincidence theorems.

To begin with we explain the notion of an H -space introduced by Horvath [6–8] and related concepts on H -spaces.

Let X be a topological space and let $\mathcal{F}(X)$ be the family of all nonempty finite subsets of X . Let $\{\Gamma_A\}$ be a family of nonempty contractible subsets of X indexed by $A \in \mathcal{F}(X)$ such that $\Gamma_A \subset \Gamma_{A'}$ whenever $A \subset A'$. The pair $(X, \{\Gamma_A\})$ is called an H -space. Given an H -space $(X, \{\Gamma_A\})$, a nonempty subset D of X is called H -convex if $\Gamma_A \subset D$ for each nonempty finite subset A of D . For a nonempty subset K of X , we define the H -convex hull of K , denoted by $H\text{-co } K$, as

$$H\text{-co } K = \bigcap \{D \subset X: D \text{ is } H\text{-convex and } D \supset K\}.$$

Then $H\text{-co } K$ is H -convex and is the smallest H -convex set containing K .

Let $(X, \{\Gamma_A\})$ be an H -space, let Y be a topological space, and let $f: X \times Y \rightarrow R$ (real number set) be a function. For each $y \in Y$, $f(x, y)$ is said to be H -quasiconvex (or H -quasiconcave) in x if the set $\{x \in X: f(x, y) < t\}$ (or $\{x \in X: f(x, y) > t\}$) is H -convex for all $t \in R$.

Now let X, Y be two topological spaces.

(1) If $A \subset X \times Y$ and $y \in Y$, let $A[y] = \{x \in X: (x, y) \in A\}$.

(2) For a multivalued mapping $T: X \rightarrow 2^Y$, we denote $T^{-1}(y) = \{x \in X: y \in T(x)\}$ and $T^*(y) = X \setminus T^{-1}(y)$ for each $y \in Y$.

(3) A function $f: X \times Y \rightarrow R$ is said to be transfer lower (upper) semicontinuous about x on X if for each $r \in R$, and each $x \in X$ and $y \in Y$, $f(x, y) > r$ ($f(x, y) < r$) implies that there exists an open neighborhood $N(x)$ of x and a point $y' \in Y$ such that $f(z, y') > r$ ($f(z, y') < r$) for all $z \in N(x)$ [17].

Remark. (a) If f is lower (upper) semicontinuous in x , then f is transfer lower (upper) semicontinuous about x on X .

(b) If function $f: X \times Y \rightarrow R$ is transfer lower (upper) semicontinuous about x on X , then the marginal function $x \rightarrow \sup_{y \in Y} f(x, y)$ is lower semicontinuous ($x \rightarrow \inf_{y \in Y} f(x, y)$ is upper semicontinuous). The converse is not true.

EXAMPLE. Let $X = Y = [0, 1]$. Suppose the function $f: X \times Y \rightarrow R$ is defined by

$$f(x, y) = \begin{cases} D(x)D(y), & \text{if } x \neq 1 \text{ and } y \neq 1. \\ 1 - D(y), & \text{if } x = 1, \\ 1 - D(x), & \text{if } y = 1, \end{cases}$$

where $D(x)$ is the Dirichlet function on $[0, 1]$. Then

$$\sup_{y \in Y} f(x, y) = 1, \quad \inf_{y \in Y} f(x, y) = 0.$$

Obviously both of the marginal functions

$$x \rightarrow \sup_{y \in Y} f(x, y)$$

and

$$x \rightarrow \inf_{y \in Y} f(x, y)$$

are continuous in X , but $f(x, y)$ is neither transfer lower semicontinuous nor transfer upper semicontinuous about x on X .

2. MAIN RESULTS

Lemmas 1 and 2 are, respectively, Theorem 2.2 and Corollary 2.3 in Tarafdar [16].

LEMMA 1. Let X be a compact topological space and let $(Y, \{\Gamma_A\})$ be an H -space. Let $T: X \rightarrow 2^Y$ be a multivalued mapping such that

- (i) For each $x \in X$, $T(x)$ is a nonempty H -convex subset of Y .
- (ii) $\{\text{int}(T^{-1}(y)): y \in Y\}$ is an open covering of X .

Then there is a continuous selection $f: X \rightarrow Y$ of T such that $f = g \circ \varphi$, where $g: \Delta_n \rightarrow Y$ and $\varphi: X \rightarrow \Delta_n$ are continuous mappings, and Δ_n is the standard n -dimensional simplex for some positive integer n .

LEMMA 2. Let $(X, \{\Gamma_A\})$ be a compact H -space and $T: X \rightarrow 2^X$ be a multivalued mapping such that

- (i) For each $x \in X$, $T(x)$ is a nonempty H -convex subset.
- (ii) $\{\text{int}(T^{-1}(y)): y \in Y\}$ is an open covering of X .

Then there is a point $x_0 \in T(x_0)$.

THEOREM 1. *Let X be a Hausdorff topological space, let $(Y, \{\Gamma_B\})$ be an H -space, and let M, N be two subsets of $X \times Y$. Suppose the following conditions are fulfilled:*

(i) *There exists a compact subset K of X such that*

(a) *for each $x \in K$, the set*

$$H - \text{co}\{y \in Y: (x, y) \notin N\} \subset \{y \in Y: (x, y) \notin M\};$$

(b) $\bigcap_{y \in Y} \text{cl}(N[y]) = \bigcap_{y \in Y} N[y]$.

(ii) *There exists a subset P of M such that P is closed in $X \times Y$, and for each $y \in Y$, the set*

$$\{x \in K: (x, y) \in P\}$$

is nonempty acyclic.

Then there exists a point $x_0 \in K$ such that $\{x_0\} \times Y \subset N$.

Proof. If the conclusion of Theorem 1 is false, then for each $x \in K$ there is a point $y_0 \in Y$ such that $(x, y_0) \notin N$. Let

$$S(x) = \{y \in Y: (x, y) \notin N\}, \quad T(x) = \{y \in Y: (x, y) \notin M\}.$$

Then $S, T: K \rightarrow 2^Y$ are such that for each $x \in K$, $S(x) \neq \emptyset$ and $H\text{-co } S(x) \subset T(x)$ by (i)(a). For each $x \in X$, there exists a point $y \in S(x)$, i.e., $x \notin N[y]$ since $S(x) \neq \emptyset$, and hence there exists a point $y' \in Y$ such that $x \notin \text{cl}(N[y'])$ by (i)(b). Consequently, there is an open neighborhood $U(x)$ of x such that $U(x) \cap N[y'] = \emptyset$. Hence $U(x) \subset S^{-1}(y')$, i.e., $x \in \text{int}(S^{-1}(y'))$. Therefore, $\{\text{int}(S^{-1}(y)): y \in Y\}$ is an open covering of K . Consequently, $\{\text{int}(H\text{-co } S)^{-1}(y): y \in Y\}$ is an open covering of K , where the mapping $H\text{-co } S: K \rightarrow 2^Y$ is defined by

$$H\text{-co } S(x) = H\text{-co}(S(x)), \quad \forall x \in K.$$

By virtue of Lemma 1 there exists a continuous mapping $f: K \rightarrow Y$ such that $f = g \circ \psi$, and

$$f(x) \in H\text{-co } S(x) \subset T(x)$$

for all $x \in K$, where $\psi: K \rightarrow \Delta_n$, $g: \Delta_n \rightarrow Y$ are continuous mappings and Δ_n is the standard n -simplex. Hence

$$(x, f(x)) \notin M, \quad \forall x \in K. \quad (1)$$

On the other hand, we define a multivalued mapping $G: Y \rightarrow 2^K$ by

$$G(y) = \{x \in K: (x, y) \in P\}, \quad \forall y \in Y.$$

By (ii), $G: Y \rightarrow 2^K$ is an upper semicontinuous multivalued mapping with nonempty closed acyclic values. Consequently, so is the mapping $F: \Delta_n \rightarrow 2^K$ defined by $F(u) = G(g(u))$. By virtue of Lemma 2.1 of [13], there exists a point $\bar{u} \in \Delta_n$ such that $\bar{u} \in \psi(F(\bar{u})) = \psi[G(g(\bar{u}))]$, and so there is a point $\bar{x} \in G(g(\bar{u})) \subset K$ such that $\bar{u} = \psi(\bar{x})$. Let $\bar{y} = g(\bar{u})$. Then $\bar{y} = g \circ \psi(\bar{x}) = f(\bar{x})$ and $\bar{x} \in G(\bar{y})$, i.e., $(\bar{x}, f(\bar{x})) = (\bar{x}, \bar{y}) \in P \subset M$. This contradicts (1) and completes the proof.

COROLLARY 2. *Let X be a Hausdorff topological space, let $(Y, \{\Gamma_B\})$ be an H -space, and let M, N be two subsets of $X \times Y$ with $M \subset N$. Suppose the following conditions are fulfilled:*

- (i) *For each $y \in Y$, the set $\{x \in X: (x, y) \in N\}$ is closed.*
- (ii) *For each $x \in X$, the set $\{y \in Y: (x, y) \notin M\}$ is H -convex or empty.*

Suppose also that there exists a subset P of M and a compact subset K of X such that P is closed in $X \times Y$ and

- (iii) *For each $y \in Y$, the set $\{x \in K: (x, y) \in P\}$ is nonempty acyclic.*

Then there exists a point $x_0 \in K$ such that $\{x_0\} \times Y \subset N$.

Remark. Both Theorem 1 and Corollary 2 improve and extend Theorem A and Theorem B to a topological space X and an H -space Y .

Theorem 3 is equivalent to Theorem 1.

THEOREM 3. *Let X be a Hausdorff topological space, let $(Y, \{\Gamma_B\})$ be an H -space, and let $F, T: Y \rightarrow 2^X$ be two multivalued mappings. Suppose the following conditions are fulfilled:*

- (i) $\bigcup_{y \in Y} F^{*-1}(y) = \bigcup_{y \in Y} \text{int}(F *^{-1}(y))$.
- (ii) $T: Y \rightarrow 2^X$ is an upper semicontinuous multivalued mapping with nonempty closed values (or T has a closed graph).
- (iii) *There exists a compact subset K of X such that*

- (a) *for each $y \in Y$, the set $T(y) \cap K$ is nonempty acyclic,*
- (b) *For each $x \in K$, $H\text{-co}(F^*(x)) \subset T^*(x)$.*

Then $\bigcap_{y \in Y} F(y) \neq \emptyset$.

Proof. Theorem 1 \Rightarrow Theorem 3: Let

$$N = \{(x, y) \in X \times Y: x \in F(y)\},$$

$$M = \{(x, y) \in X \times Y: x \in T(y)\}.$$

Then for each $x \in K$, the set

$$\begin{aligned} H\text{-co}\{y \in Y: (x, y) \notin N\} &= H\text{-co}(Y \setminus F^{-1}(x)) \\ &\subset Y \setminus T^{-1}(x) \\ &= \{y \in Y: (x, y) \notin M\} \end{aligned}$$

by (iii)(b). By (i) $\bigcap_{y \in Y} cl(N[y]) = \bigcap_{y \in Y} N[y]$, by (ii) M is closed, and by (iii)(a) the set

$$\{x \in K: (x, y) \in M\} = K \cap T(y)$$

is nonempty acyclic for each $y \in Y$.

By Theorem 1, there exists a point $x_0 \in K$ such that $\{x_0\} \times Y \subset N$, i.e., $(x_0, y) \in N$ for all $y \in Y$. Hence $x_0 \in \bigcap_{y \in Y} F(y) \neq \emptyset$. This completes the proof.

Theorem 3 \Rightarrow Theorem 1: For each $y \in Y$, let

$$S(y) = \{x \in X: (x, y) \in M\},$$

$$T(y) = \{x \in X: (x, y) \in P\},$$

$$F(y) = \{x \in X: (x, y) \in N\}.$$

By (i) there exists a compact subset $K \subset X$ such that $K \cap T(y)$ is nonempty acyclic and for each $x \in K$ the following hold:

$$(a) \quad H\text{-co}(F^*(x)) \subset S^*(x) \subset T^*(x).$$

$$(b) \quad \bigcup_{y \in Y} \text{int}(F^*{}^{-1}(y)) = \bigcup_{y \in Y} F^*{}^{-1}(y).$$

Since again P is closed in $X \times Y$, T has a closed graph. Consequently by Theorem 3 $\bigcap_{y \in Y} F(y) \neq \emptyset$, and hence there is a point $x_0 \in \bigcap_{y \in Y} F(y)$. Thus $\{x_0\} \times Y \subset N$ and Theorem 1 is proved.

COROLLARY 4. Let X be a Hausdorff topological space, let $(Y, \{\Gamma_B\})$ be an H -space, and let $F, T: Y \rightarrow 2^X$ be two multivalued mappings. Suppose the following conditions are fulfilled:

$$(i) \quad \bigcup_{y \in Y} F^*{}^{-1}(y) = \bigcup_{y \in Y} \text{int}(F^*{}^{-1}(y)).$$

(ii) $T: Y \rightarrow 2^X$ is an upper semicontinuous multivalued mapping with nonempty closed values (or T has a closed graph),

(iii) *There exists a compact subset K of X such that*

(a) *for each $y \in Y$, the set $T(y) \cap K$ is nonempty acyclic and $T(y) \subset F(y)$,*

(b) *for each finite subset $\{y_1, y_2, \dots, y_n\} \subset Y$,*

$$T(\Gamma_{\{y_1, y_2, \dots, y_n\}}) \subset \bigcup_{i=1}^n T(y_i).$$

Then $\bigcap_{y \in Y} F(y) \neq \emptyset$.

Proof. Since $T(y) \subset F(y)$ for each $y \in Y$, $T^{-1}(x) \subset F^{-1}(x)$ for each $x \in X$, and hence $F^*(x) \subset T^*(x)$ for each $x \in X$. Hence, it is sufficient to show that $T^*(x)$ is H -convex. For each finite subset $\{y_1, y_2, \dots, y_n\} \subset T^*(x)$, we have $x \notin \bigcup_{i=1}^n T(y_i)$, and hence $x \notin T(\Gamma_{\{y_1, y_2, \dots, y_n\}})$. Therefore, for each $y \in \Gamma_{\{y_1, y_2, \dots, y_n\}}$, we have $x \notin T(y)$, i.e., $y \in T^*(x)$. Hence $\Gamma_{\{y_1, y_2, \dots, y_n\}} \subset T^*(x)$. This shows that $T^*(x)$ is H -convex and completes the proof.

Remark. Even if X and Y are nonempty convex sets of topological vector spaces and $F = T$, Theorem 3 and Corollary 4 are different from Theorems 2.4 and 2.5 in [19], since our multivalued mapping is of upper semicontinuous type.

THEOREM 5. *Let X be a compact Hausdorff topological space, let $(Y, \{\Gamma_A\})$ be an H -space, and let $f, g: X \times Y \rightarrow R$ be two functions. Let $\beta = \sup_{x \in X} \inf_{y \in Y} g(x, y)$ and $D = \{(x, y) \in X \times Y: f(x, y) > \beta\}$. Suppose that the following conditions are fulfilled:*

(i) *$f(x, y) \leq g(x, y)$ for all $(x, y) \in D$.*

(ii) *$g(x, y)$ is transfer upper semicontinuous in x , and for each $y \in Y$, the set $\{x \in X: f(x, y) > t\}$ is acyclic or empty for each $t > \beta$.*

(iii) *For each $x \in X$, $f(x, y)$ is H -quasiconvex in y , and $f: X \times Y \rightarrow R$ is upper semicontinuous. Then*

$$\inf_{y \in Y} \sup_{x \in X} f(x, y) \leq \sup_{x \in X} \inf_{y \in Y} g(x, y).$$

Proof. If the conclusion of Theorem 5 is false, then there is a real number t such that

$$\beta = \sup_{x \in X} \inf_{y \in Y} g(x, y) < t < \inf_{y \in Y} \sup_{x \in X} f(x, y).$$

For each $y \in Y$, let $F(y) = \{x \in X: g(x, y) \geq t\}$, $T(y) = \{x \in X: f(x, y) \geq t\}$. Then $F, T: Y \rightarrow 2^X$ are two multivalued mappings with $T(y) \subset F(y)$ for all $y \in Y$, since $f(x, y) \leq g(x, y)$ for all $(x, y) \in D$.

Since f is upper semicontinuous, the graph $\text{Gr}(T)$ of T is closed, and since X is compact, $T: Y \rightarrow 2^X$ is upper semicontinuous. Since again, for each $x \in X$, $f(x, \cdot)$ is H -quasiconvex, the set

$$T^*(x) = \{y \in Y: x \notin T(y)\} = \{y \in Y: f(x, y) < t\}$$

is H -convex.

For each $x \in \bigcup_{y \in Y} F^{*-1}(y)$ there is a point $y_0 \in Y$ such that $g(x, y_0) < t$. By the first part of condition (ii), there exists an open neighborhood $U(x)$ of x and a point $y' \in Y$ such that $g(z, y') < t$ for all $z \in U(x)$, and hence $U(x) \cap F(y') = \emptyset$. Hence $x \in \text{int}(F^{*-1}(y'))$. It shows

$$\bigcup_{y \in Y} F^{*-1}(y) = \bigcup_{y \in Y} \text{int}(F^{*-1}(y)).$$

Note that since X is compact by the second part of (ii) and (iii) for each $y \in Y$, the set

$$T(y) = \{x \in X: f(x, y) \geq t\} = \bigcap_{0 < \varepsilon < t - \beta} \{x \in X: f(x, y) > t - \varepsilon\}$$

is acyclic (this follows from the continuity of Čech homology).

Therefore all conditions of Theorem 3 are fulfilled, so by Theorem 3, $\bigcap_{y \in Y} F(y) \neq \emptyset$. Thus there exists a point $x_0 \in X$ such that $x_0 \in \bigcap_{y \in Y} F(y)$, i.e., $g(x_0, y) \geq t$ for all $y \in Y$. Hence

$$\sup_{x \in X} \inf_{y \in Y} g(x, y) \geq t.$$

This contradicts the choice of t and completes the proof.

COROLLARY 6. *Let X be a compact Hausdorff topological space, and let $(Y, \{\Gamma_A\})$ be an H -space. If an upper semicontinuous function $f: X \times Y \rightarrow R$ is such that*

(i) *For each $y \in Y$, the set $\{x \in X: f(x, y) > t\}$ is acyclic or empty for each $t \in R$,*

(ii) *For each $x \in X$, $f(x, y)$ is H -quasiconvex in y , then*

$$\inf_{y \in Y} \max_{x \in X} f(x, y) = \max_{x \in X} \inf_{y \in Y} f(x, y).$$

Proof. By Theorem 5 we have

$$\inf_{y \in Y} \sup_{x \in X} f(x, y) = \sup_{x \in X} \inf_{y \in Y} f(x, y).$$

Note that X is compact and f is upper semicontinuous, so we have

$$\inf_{y \in Y} \max_{x \in X} f(x, y) = \max_{x \in X} \inf_{y \in Y} f(x, y).$$

This completes the proof.

Remark. In the case where X is compact, Corollary 6 contains Theorem 4 of Ha [5] as a special case.

THEOREM 7. *Let K be a nonempty compact subset of a Hausdorff topological space X , and let $(Y, \{T_A\})$ be an H -space. Let $T: X \rightarrow 2^Y$ and $F: Y \rightarrow 2^X$ be multivalued mappings such that*

- (i) F has a closed graph $\text{Gr}(F)$.
- (ii) $\{\text{int}(T^{-1}(y)); y \in Y\}$ is an open covering of X .
- (iii) For each $x \in K$, $T(x)$ is nonempty H -convex.
- (iv) For each $y \in Y$, $F(y) \cap K$ is nonempty acyclic.

Then there exist a point $x_0 \in X$ and a point $y_0 \in Y$ such that $x_0 \in F(y_0)$ and $y_0 \in T(x_0)$.

Proof. For each $x \in X$, let $G(x) = F^{-1}(x) = \{y \in Y: x \in F(y)\}$. Then by (i) $G: X \rightarrow 2^Y$ has a closed graph and by (iv) for each $y \in Y$,

$$\begin{aligned} G^{-1}(y) \cap K &= \{x \in X: y \in G(x)\} \cap K \\ &= \{x \in X: x \in F(y)\} \cap K \\ &= F(y) \cap K \end{aligned}$$

is nonempty acyclic. Let $P = \text{Gr}(G)$ and $M = N = \{(x, y) \in X \times Y: y \notin T(x)\}$. Then for each $y \in Y$, the set

$$\{x \in K: (x, y) \in P\} = \{x \in K: x \in G^{-1}(y)\} = K \cap G^{-1}(y)$$

is nonempty acyclic, and by (iii) P is closed in $X \times Y$. If the conclusion of Theorem 7 is false, then $T(x) \cap G(x) = \emptyset$ for all $x \in X$, and hence $P \subset M$. Moreover, by (iii) we know that for each $x \in K$, the set

$$\{y \in Y: (x, y) \notin M\} = T(x)$$

is H -convex, and by (ii) $\bigcap_{y \in Y} \text{cl}(N[y]) = \bigcap_{y \in Y} N[y]$.

So by Theorem 1, there exists a point $x_0 \in K$ such that $\{x_0\} \times Y \subset M$, i.e., $y \notin T(x_0)$ for all $y \in Y$. Hence $T(x_0) = \emptyset$. This contradicts (iii) and completes the proof.

Remark. Theorem 7 improves Theorem 3.1 of [2], and improves and generalizes Theorem 2.3 of [12], Theorem 7 of [15], Theorem 1 of [9], Theorem 1 of [1], and Theorem 1 of [18].

REFERENCES

1. F. E. Browder, Coincidence theorems, minimax theorems, and variational inequalities, *Contemp. Math.* **26**, (1984), 47–80.
2. X. P. Ding and E. Tarafdar, Some coincidence theorems and applications, *Bull. Austral. Math. Soc.* **50** (1994), 73–80.
3. K. Fan, A generalization of Tychonoff's fixed point theorem, *Math. Ann.* **142** (1961), 305–310.
4. K. Fan, Applications of a theorem concerning sets with convex sections, *Math. Ann.* **163** (1966), 189–203.
5. C. Ha, Minimax and fixed point theorems, *Math. Ann.*, **248**, (1980), 73–77.
6. C. Horvath, Point fixes et coïncidences pour les applications multivoques sans convexité, *C. R. Acad. Sci. Paris* **296** (1983), 403–406.
7. C. Horvath, Point fixes et coïncidences dans les espaces topologiques compacts contractiles, *C. R. Acad. Sci. Paris* **299** (1984), 519–521.
8. C. Horvath, Some results on multivalued mappings and inequalities without convexity, in "Nonlinear and Convex Analysis" (B. L. Lin and S. Simons, Eds.), pp. 99–106, Marcel Dekker, New York, 1987.
9. H. Kormiya, Coincidence theorems and saddle point, *Proc. Amer. Math. Soc.* **96** (1986), 59–62.
10. F. C. Liu, On a form of KKM principle and SupInfSup inequalities of von Neumann and of Ky Fan Type, *J. Math. Anal. Appl.* **155** (1991), 420–437.
11. J. F. McClendon, Minimax and variational inequalities for compact spaces, *Proc. Amer. Math. Soc.* **89**(4) (1983), 717–721.
12. G. Mehta and S. Sessa, Coincidence theorems and maximal elements in topological vector spaces, *Math. Japon.* **47** (1992), 839–845.
13. S. Park, J. S. Bae, and H. K. Kang, Geometric properties minimax inequalities, and fixed point theorems on convex spaces, *Proc. Amer. Math. Soc.* **121**(2) (1994), 429–439.
14. M. Sion, On general minimax theorems, *Pacific J. Math.* **8** (1958), 171–176.
15. S. Sessa, Some remarks and applications of an extension of a lemma of Ky Fan, comment, *Math. Univ. Carolin.* **29**, (1988), 567–575.
16. E. Tarafdar, Fixed point theorems in H -spaces and equilibrium points of abstract economies, *J. Austral. Math. Soc.* **53** (1992), 252–260.
17. G. Q. Tian, Generalizations of the FKKM theorem and the Ky Fan minimax inequality, and applications to maximal elements, price equilibrium and complementarity, *J. Math. Anal. Appl.*, **170** (1992), 457–471.
18. E. Tarafdar, On nonlinear variational inequalities, *Proc. Amer. Math. Soc.* **67** (1977), 95–98.
19. G. H. Greco, Minimax theorems and saddling transformations, *J. Math. Anal. Appl.* **147** (1990), 180–197.